

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



23 JUL 2004



(43) International Publication Date  
21 August 2003 (21.08.2003)

PCT

(10) International Publication Number  
**WO 03/068951 A1**

(51) International Patent Classification<sup>7</sup>: **C12N 9/80,**  
15/57, A01K 67/027, A61K 38/50

(21) International Application Number: **PCT/SG02/00011**

(22) International Filing Date: 23 January 2002 (23.01.2002)

(25) Filing Language: English

(26) Publication Language: English

(71) Applicant (for all designated States except US): **INSTITUTE OF MOLECULAR AGROBIOLOGY [SG/SG];**  
1 Research Link, National University of Singapore, Singapore 117604 (SG).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **ZHANG, Lian, Hui [AU/SG];** 360 Pasir Panjang Road., #03-11 Goldcoast Condominium, Singapore 118699 (SG). **LIN, Yi, Han**

[AU/SG]; 360 Pasir Panjang Road, #04-10 Goldcoast Condominium, 118699 Singapore (SG). **XU, Jin, Ling** [AU/SG]; 360 Pasir Panjang Road, #03-11 Goldcoast Condominium, Singapore 118699 (SG).

(74) Agent: **ELLA CHEONG MIRANDAH & SPRUSONS;** Robinson Road Post Office, P.O. Box 1531, Singapore 903031 (SG).



(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),

[Continued on next page]

(54) Title: **RALSTONIA AHL-ACYLASE GENE**

(57) Abstract: This invention provides a gene, *qsba*, which encodes a protein useful for inactivating certain bacterial quorumsensing signal molecules (N-acyl homoserine lactones) which participate in bacterial virulence and biofilm differentiation pathways. This gene was isolated from *Ralstonia sp.*, strain XJ12B. The invention also provides the Qsba protein, which possesses N-acyl homoserine lactone inactivating activity.



WO 03/068951 A1

BEST AVAILABLE COPY



Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),  
European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR,  
GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent  
(BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR,  
NE, SN, TD, TG).

*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

**Published:**

— *with international search report*

## RALSTONIA AHL-ACYLASE GENE

## BACKGROUND OF THE INVENTION

Field of the Invention

[0001] This invention pertains to the field of molecular biology. In particular, the invention relates to an N-acyl homoserine lactone acylase gene from *Ralstonia* sp. XJ12B.

Description of the Background Art

[0002] N-acyl homoserine lactones (AHLs), also known as autoinducers, are widely used quorum sensing signal molecules in many Gram-negative bacteria. These compounds regulate certain classes of target genes in bacteria, such as virulence genes or biofilm differentiation genes. Generally, quorum sensing molecules are highly conserved and share an identical homoserine lactone moiety. The length and structure of their acyl side chains are different, however. Although the target genes regulated by AHLs in different bacteria species are varied, basic mechanisms of AHL biosynthesis and gene regulation are conserved among different bacterial species.

[0003] The general feature of AHL-mediated gene regulation is that it is cell population dependent (quorum sensing). Bacteria secrete AHLs into the environment; extracellular concentration of AHLs increases as bacterial cell populations grow. When AHL accumulates to a threshold extracellular concentration, the expression of certain sets of target genes are triggered in the bacteria.

[0004] Bacteria using these signals release, detect and respond to the accumulation of AHL signal molecules for synchronizing expression of a particular sets of genes and coordinating cellular activities within the bacterial cell population. AHLs are involved in regulation of a range of biological functions, including bioluminescence in *Vibrio* species (13, 4), Ti plasmid

conjugal transfer in *Agrobacterium tumefaciens* (31), induction of virulence genes in *Burkholderia cepacia*, *Erwinia carotovora*, *Erw. chrysanthemi*, *Erw. stewartii*, *Pseudomonas aeruginosa*, and *Xenorhabdus nematophilus* (3, 6, 12, 17, 19, 22, 23, 24, 26), regulation of antibiotic production in *P. aureofaciens* and *Erw. carotovora* (6, 26), swarming motility in *Serratia liquifaciens* (14) and biofilm formation in *P. fluorescens* and *P. aeruginosa* (1, 8). In many other bacterial species the relevant biological functions controlled by AHLs remain to be investigated (2, 5, 11).

[0005] A number of plant, animal and human bacterial pathogens use AHL quorum-sensing signals to regulate expression of pathogenic genes and aid in the formation of biofilms. Therefore, AHL quorum-sensing signal molecules are group of molecular targets for genetic and chemical manipulations since disruption of these signaling mechanisms can prevent or reduce the ability of these bacteria to infect plant and animal tissues or to form biofilms.

[0006] The gene encoding an AHL-inactivation enzyme (AiiA) from a Gram-positive bacterium (*Bacillus* strain 240B1) has been cloned (9). AiiA (also known as AHL-lactonase) inactivates AHL activity by hydrolyzing the lactone bond of AHLs (10). Expression of *aiiA* in transformed *Erw. carotovora* (a pathogenic strain which causes soft rot disease in many plants) significantly reduces the release of AHL, decreases extracellular pectrolytic enzyme activities, and attenuates pathogenicity on potato, eggplant, Chinese cabbage, carrot, celery, cauliflower, and tobacco (9). Transgenic plants expressing AHL-lactonase showed a significantly enhanced resistance to *Erw. carotovora* infection and delayed development of soft rot symptoms (10). AHL-inactivation mechanisms appear to be widely distributed. For example, a bacterial isolate of *Variovorax paradoxus* was reported to use AHL molecules as its energy and nitrogen sources, indicating the possible presence of AHL-degrading enzymes (18).

[0007] Further methods to counteract AHL-mediated plant, animal and human disease and plant pathogen virulence by interfering with bacterial intercellular communication would be highly desirable.

#### SUMMARY OF THE INVENTION

[0008] Accordingly, in this study, the cloning and characterization of a gene encoding an AHL-acylase from a bacterial isolate *Ralstonia* sp. JX12B is reported.

[0009] In one embodiment, the invention provides a composition of matter which comprises a nucleic acid according to SEQ ID NO: 1. In another embodiment, the invention provides a composition of matter which comprises a nucleic acid selected from the group consisting of nucleotides 1234-3618 of SEQ ID NO: 1, a fragment thereof and a substantially homologous variant thereof.

[00010] In yet a further embodiment, the invention provides a nucleic acid according to claim 2 which comprises nucleotides 1234-3618 of SEQ ID NO: 1.

[00011] In yet a further embodiment, the invention provides a composition of matter which comprises a peptidic sequence selected from the group consisting of a peptidic sequence according to SEQ ID NO: 2, a fragment thereof and a substantially homologous variant thereof.

[00012] In yet a further embodiment, the invention provides a composition of matter which comprises a peptidic sequence encoded by a nucleic acid selected from the group consisting of nucleotides 1234-3618 of SEQ ID NO: 1, a fragment thereof and a substantially homologous variant thereof.

[00013] In yet a further embodiment, the invention provides a composition of matter which comprises a peptidic sequence selected from the group consisting of SEQ ID NO: 2, a fragment thereof, a subunit thereof and a substantially homologous variant thereof, such as a peptidic sequence according to SEQ ID NO: 2, a peptidic sequence comprising amino acids 36-217 233-794[?] of SEQ ID NO: 2 or a peptidic sequence comprising amino acids 233-794 of SEQ ID NO: 2.

[00014] In yet a further embodiment, the invention provides a composition of matter as described above which inactivates AHL.

[00015] In yet a further embodiment, the invention provides a method of modulating AHL signaling activity which comprises contacting said AHL with a composition of matter as described above.

[00016] In yet a further embodiment, the invention provides a transgenic plant or non-human mammal harboring a nucleic acid as described above.

[00017] In yet a further embodiment, the invention provides a method of controlling a bacterial disease in a mammal which comprises administering to said mammal a composition of matter as described above, wherein the expression of pathogenic genes of said bacteria are regulated by AHL signals.

[00018] In yet a further embodiment, the invention provides a method of controlling a bacterial disease in a plant which comprises administering to said plant a composition of matter as described above, wherein the expression of pathogenic genes of said bacteria are regulated by AHL signals.

#### BRIEF DESCRIPTION OF THE DRAWINGS

[00019] Figure 1 is a photograph showing AHL inactivation bioassay results for bacterial cultures and bacterial proteins from the indicated bacterial clones. Figure 1A shows the results of a bioassay with bacterial cultures of *E. coli* DH5 $\alpha$  strains 13H10 (slice 1), 2B10 (slice 2), MUB3 (slice 3), MUC6 (slice 4), GST-QsbA (slice 5) and GST (slice 6), which contain plasmid clones or constructs p13H10, p2B10, pMUB3, pMUC6, pGST-QsbA, and pGST, respectively. Figure 1B shows results for bioassay of the indicated bacterial proteins GST-QsbA and GST.

[00020] Figure 2 is a graph showing the temperature and pH optimum profiles of AHL acylase.

[00021] Figure 3 is a graph showing the time course of OOHL inactivation by the purified AHL-acylase.

## DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

[00022] A bacterial isolate of *Ralstonia* sp. XJ12B from a biofilm sample in a water treatment system was found to enzymatically inactivate AHLs, bacterial quorum-sensing molecules, in a bioassay using *Agrobacterium tumefaciens* strain Nt1 (*traR*; *tra::lacZ749*) as an indicator for AHL activity. The gene encoding the protein exhibiting this enzyme activity for AHL inactivation (*qsba*) was cloned from a bacterial strain isolated from the biofilm sample and found to encode a peptide of 794 amino acids.

[00023] Bacterial cultures and bacterial proteins were assayed for the ability to inactivate AHL using *Agrobacterium tumefaciens* indicator cells. A *tumefaciens* was cultured at 28°C in MM medium as described in Zhang et al. (31). The bacteria or protein to be assayed is first mixed with an AHL substrate, for example N- $\beta$ -oxooctanoyl-L-homoserine lactone (OOHL), and the reaction (inactivation of the AHL) is allowed to proceed. If AHL inactivation activity is present in the sample (i.e. the AHL has been cleaved and inactivated), then the inactivated AHL products fail to trigger the expression of *lacZ* reporter gene which is under the control of a TraR-dependent promoter. The strain A. *Tumefaciens* NT1 hosting the *lacZ* reporter system therefore does not turn blue in the presence of substrate 5-bromo-4-chloro-3-indolyl  $\beta$ -D-galactopyranoside (X-gal). See Example 2 for details of the bioassay. Any AHL may be used in the assay, as desired. Of course, any suitable assay for cleavage of AHL, including traditional *in vitro* enzyme assays may be used to detect the AHL inactivation activity. Those of skill in the art are able to modify or devise assays to detect and/or quantitate AHL inactivation.

[00024] *Escherichia coli* strain DH5a was used as a host for DNA manipulation. Both *Ralstonia* sp. and *E. coli* were cultured in LB medium (tryptone, 10 g/L, yeast extract, 5 g/L, and NaCl, 10 g/L, pH 7.0) at 37°C. Appropriate antibiotics were added when necessary at the following concentrations: ampicillin, 100  $\mu$ g/ml; tetracycline, 10  $\mu$ g/ml; and kanamycin, 20  $\mu$ g/ml.

**[00025]** The gene encoding the protein responsible for the detected AHL inactivation was isolated using a cosmid library of 1600 clones with the genomic DNA of *Ralstonia* sp. strain XJ12B, constructed in *E. coli*. *E. coli* transfectants were screened for the ability to inactivate AHL. One clone, p13H10, was found to inactivate AHL. Cosmid DNA from p13H10 was digested, fused into a cloning vector, ligated and transformed into *E. coli*. The *E. coli* clones again were assayed for AHL inactivating activity. One clone, containing a 4 kb insert, had AHL inactivation activity.

**[00026]** Plasmids were subsequently purified for sequencing. The 4 kb fragment from clone p2B10 was completely sequenced according to known methods using ABI Prism dRhodamine Terminator Cycle Sequencing Ready Reaction Kit (Perkin-Elmer Applied Biosystems). See Table I, below. The sequence contained an open reading frame of 2385 nucleotides which was the AHL inactivation gene, *qsba*, encoding a predicted polypeptide of 794 amino acids (85,373 Daltons).



**Table I. *QsbA* Gene (*Ralstonia* sp.) Nucleotide Sequence (SEQ ID NO: 1).**

```

gtttgggaagtgggagcgcgctgtgcag  cgcggcgccctcagcgcgcgagctcgccg  cgcaccgaatgcgcgcgcgggtggcgccc  90
ggcggtggcggtgtggcgcgcgatcagg  cgccggaaggcgacatgtctgataaccg  cactgttcggcgattgcccgcaggtcagc  180
gtgctgacttccagcaggtggcaggcgcg  tccacgcgcagcggtgcagcaattgcagc  ggcgaggtgccagggtcttgggtgaatgc  270
cgcagcagcgtgcgctcgtggtcgaggcg  gcggcgccagcttggccaggtcgaacggc  tcgtgcaggtgctgctgcaggtagcgccg  360
gccccgagtaccacgctggtgcggtggcg  ggcttgcctgcgcagcgagatggcggtgg  tcaccgcgcgacgggtggtcgagcagggc  450
tgccgaggggtgcgtgccagcggtgtcg  gccagggcgccgaccaggcgctgcgtgagc  gccacgcgctgtccatcgcgcgccgctc  540
agcacgttgccgctgctgacgatggcctgc  tccgccaccaccttcagctgcgggtagttg  ccgtgcagcagcgccgcatcagccagtc  630
accgtcaagcgccggcgccggcgagcgcg  ccggccagcagcgcccgccgggtgaaggac  gaggccaccaagcctgcccgcgctccaggt  720
gcgcggatggtggcgcttcccactccag  caggggcaggcgctgctccagcgctgat  gtggtcgaaatgcaggggcgggacgaccag  810
cgcgctgccagcgcgcgctgcggcgcg  cagcggtgcgcagcgcgagggcagggtctc  ggcgggcgccctgccagcgggcggtcgcg  900
cgcgaccagcgccaccggaacacgggct  ggcgcatcgcgacgcttggccagcgagggt  ggagggcgagcgattggccagcgagggt  990
gtcgcgacgggtgcggcggtggagggcc  ggcgtcgggaaaggtcagcaggtcgatgtc  ggcacccgaaagtataggggagcgggcg  1080
gagggcctcctgctggcgggattgacccca  actctggcggaataaccttctcctccggg  cgggccccagtcgacgatacggcggtggct  1170
gcgctgcgcgcgcgcgcaagactagagcg  acacaagacaagaccgacagggagacaa  cgcATGATGCAGGGATTTCGCGTGGCGGC  1260
ACGCTCGCCATGGCCGCGCTCGCGGCGCTG  GCCGCTGCGCCAGTTCACCGATGGCGCG  TGGGGTTCGCTCAGCGACACCGGCTGTCC  1350
GCCGAGATCCGCGCAGCGGCTTCGGCATT  CCGCACATCCGCGCAACGACTACGCCAG  CTCGGCTATGGCATGGCCTATGCCTACGCG  1440
CAGGACAACCTTGCTGCTGGCCGACACAG  GTGGTCACCGTCAACGGCGAGGCGCTCGA  ACCTTCGGGCCCCGAGGGCAGCGTGCAGCG  1530
TCGTTCAAGCCGATCCCCAACCTGCAGTCG  GACGCCCTTCCTCAAGGGCATCTTCGACG  GACGGCTTCGCGCGCGGTTATGCGCAGATG  1620
TCGCGGAGGCGCGGAGTGTGCGCGCGG  TACATCGCGCGCTTCAACCGCTATCTCAA  GACACGCCGCGCCGCAACTTCCCGGCGGCC  1710
TGCCGCAATGCCGCTTGGTGTGCTCGCTC  ACGCTCAGCGACATGATGCGCATGGGCGA  GAGAAGGCGATCCAGGCCAGCGCCGCGGCC  1800
ATGCTGGCGGGCATCGTGCAGCGCGAGCC  CCGGGCGCGACGCCGGTGGCCGAGCGCGAG  ATTCGCGCGCAGGCGCTCGACACCGTGGCG  1890
CTGACCGCGAATCGCACTTCCCGTGGACG  CCGATCGGCTCCAACGCGTGGGCGCTTCG  GCTGACGCCACCGCAACCGGCGCGCGTG  1980
GCCTCGATCGCGGCTTCCCGGTGGTGAGC  ACCACCAACCGTTCTTACAGGTCCACCTG  ACGGTGCCCGGCAAGCTCGAGCTGAGGCG  2070
TTGTTCAAGTGAAGTGGCGGAAGCGGAC  ATCGGCTTCAACAAGGACGTGGCGTGGAC  CACACCGTCTCCACCGCGCGCGCTTCAAC  2160
GACGTGAAGTGGCGGACGGCGCTCGAG  CCGACACCTACTGGTTCGACGGCAGCGCG  CACAAGATGACCACCGCAGGTCGCGCTTC  2250
GGCATGCCGTGGACACGAGAGGCGCTAC  CGCGCACGACACCTTCTACGACACCATC  TACGGCCCGGTGCTGTGATGCCGAGCGGC  2340
CAGGCCCGGACGTTGGCGGCGATCCGCCAG  GCCATCGGCAACTGGGCTTCCCTGGGTC  CGCTCGGTTCGACAGTGGCTGCATATCGGG  2430
CGCGTGTTCGCGGACGTGCGACACGCGG  GACGTGCCGGCGCGGAGCTCCAGCGCTGT  AACACCATCGCCACCGACCGCAACGCGCG  2520
AAGACGCGGCGCTGGTGTGCTCGACGGC  TCGCGCGGACCTGCACTGGCAGGTGCGAT  GCCCGTTCGCGGTTACCGGGCTGGTGGCG  2610
CCC GCGCGATGCCGGTGTGCGAGCGGAC  GACTACGTGCGCAACAGCAATGACAGTCC  CCGGCTTCGCGGTTACCGGGCTGGTGGCG  2700
GGCTTCTCGCGGTGATGGGCTCGGTCGAC  GTACCGAGCGGCTGCGCACGCGATCGGC  TGGCTGACCAACCCCGCGCAAAAGCTGACC  2790
ACCGAGGACTGCCCGGCAACCGCATCGAT  CTGCCGAACTGCAGGCGATGATCTTACG  CTGATCGAGATCGGCCGCGCTGGCGCGC  2880
GGCGACCTGCTCGCGGCGATGCAAGGCCAG  CCGGCCCGGATGCCGAGTGGCGGACGGC  AATGCCAACCTGGCGGGACAACTGGTGTG  2970
AGCAACGCGACGCGCGCGCGCGCACTG  TTCCGCGAGTTCTGAGTGGCGGCAAGGAC  TCGCGCGCCTCGGCCAGTGGAAACCGCAC  3060
CCGCGGACCGCGTCCACACGCGCGCGCGG  CTGCGCATGAACGACGCGAGGTACGCACG  ATCGCGCAGGTGCAGCGCTCGATTCGAC  3150
GGCGCGGTGCGCAAGCGCGGCTTCGCGCTG  GATGCGCGCTGGGACGGTACAGGCGCGG  GCGGTGTTCAAGGCGCTGAAGGAAGCGTG  3240
GGCGCGGAGGAATACGAAGCGCTGCTCAAC  AAGCTGCAAACTGCGCATCGGGCCGAG  CACGCACCGGACGGCTCCATCGCCCTGCAC  3330
ATCCAGACCGTGACCTTCGACGACCGGGC  CCGGTGCGCGACGCCATCTCACCTACGGC  GGGCTGCCGCTGTATTTGCGCACAGCTAC  3420
TTCGACAGATGCGTGCCTACTCGGGCAAG  CACTGGAACCGGCTGCGGTTCTCGGAAGC  GAATCGACCGACACGCGCTCGCCGACGCG  3510
ATGCGGTTGTCGAGTGagggtgcccgtg  cctggaataacccccgcttggcgggcg  GCCATCGCGGCGGATCCGGCGCTGAAGGTG  3600
gttggaaacgcacccgacatgactgtat  tgtgactctgctgtgtccgtgt 3743  ttttttgcagtgatgaatggctcaatcgt  3690

```

The predicted open reading frame of the *qsbA* gene is shown in upper case letters with the start codon and stop codon in bold. A putative ribosome binding site (AGGAGA) is underlined.

**[00027]** Sequence analysis of this peptide indicated that *QsbA* did not have any significant homology with the known AHL-lactonase quorum-sensing molecule inactivator encoded by the *aiiA* gene from *Bacillus* sp. 240B1, however the deduced peptide sequence was typical of the primary structure of aculeacin A acylases (AACs) and penicillin G acylases, with signal peptide- $\alpha$  subunit-spacer- $\beta$  subunit organization (16, 30). The *Ralstonia* sequence shares substantial identity with AACs from *Deinococcus radiodurans* strain R1, *Actinoplanes utahensis* and a putative acylase from *Pseudomonas aeruginosa*, all of which catalyze deacylation of their substrates. These AAC genes are translated as single precursor polypeptide and then processed to the active

form, which has two subunits. Aculeacin A is an echinocandin-type antifungal antibiotic with a long fatty side chain. Aculeacin A acylases purified from *A. utahensis* catalyze the hydrolysis of the amide bond on the palmitoyl side chain of aculeacin A (29). The primary structure of the protein, as well as enzyme activity analysis with different substrates, discussed below, therefore indicates that *qsbA* encodes an AHL-acylase which cleaves the amide linkage between the acyl side chain and the homoserine lactone moiety of AHLs.

[00028] The presumed  $\alpha$  and  $\beta$  subunits of QsbA are located at amino acid positions 36-217 and 233-794, respectively, of SEQ ID NO: 2, with a 15 amino acid spacer between them, as determined by alignment with the peptide sequences from *D. radiodurans* strain R1, *A. utahensis* and *P. Aeruginosa*. See Table II.

Table II. Aligned Amino Acid Sequences of *QsbA* from *Ralstonia* sp. XJ12B (SEQ ID NO: 2), *D. radiodurans* strain R1 acylase (SEQ ID NO: 3), *A. utahensis* acylase (SEQ ID NO: 4) and *P. aeruginosa* acylase (SEQ ID NO: 5).

R.sp	MMQGF---ALRGTLAMAALAALAGCA-----SSTDGRWGSLSDTGLSAEIRRTGFGIPHIRANDYASLGYGMAAYAYAQDN	72
D.rad	MSR-----SPFSSVSLPARLLLSLL-----LGPLMLGGAASAQTYQVQIQRTAHGIPHIQASDLGGIGYGVGYSYAQDN	70
A.uta	MTSSY---MRLKAAAIAGFVIVATAA-----VPSPAS-GREHDGGYAAIIRRASYGVPHITADDFGSLGFGVGYVQAEDN	71
P.aer	MSRFFRPPLCRETTSMGMRVTLTGLAGMLLGSMMPVQADMPRPTGLAADIRWTAYGVPHIRAKDERGLGYGIGYAYARDN	80
	* : : : . * : : : * * . : : * . * . *	
	1	
R.sp	LCLLADQVVTNNGERSKTFGPEGTVTVSFKPIPNLQSDAFFKGFDEEDGLRAGYAQMSPEARELLRGYIAGFNRYLKDTP	152
D.rad	LCLLADQVMTVRGERSKFLGAEGKTVVGFQPVNLLSDVFFKTVIEPGRLOAGYRDQ-PQILALMRGYVAGVNRYLKDTTP	149
A.uta	ICVIAESVVTANGERSRWFGATGPDADVRTTSSTQAIDDRVAERLLEGPDRDGVAPCDDVDRDQMRGFVAGYNHFLRRTG	151
P.aer	ACLAEIIVTARGERARYFGSEGKSSAELD---NLPSDIFYAWLNQPEALQAFWQAQTPAVRQLLEGYAAGFNRLREAD	157
	* : : : * . * . . . :	
R.sp	PANFP-AACRNAAWVRPLTLGDMMRMGEEKAIQASAGAMLAGIVAAQPPGRTFVAEREIPPQAVDTVALDRELQLRDMPI	231
D.rad	PEQWP-SACRNADWVRPLTELDMVRLGEEKAIQASAGAMVSAITSARPPQ---AGASTAAPRPDLAAFNRQYRFNDLPI	224
A.uta	VHRLTDPACRGKAWVRPLSEIDLWRTSWDSMVBRAGSGALLDGIVAATPPT---AAGPASAPEAPDAAAIAAALDGTSGAGI	228
P.aer	GKTTS---CLGQFWLRAIATDDLRLRLTRLLVEGGVGFADALVAAAPPG---AEKVALSGEQAFQVAEQRRQRFLER	230
	. * . * : : * : * : . . . * : : : * * *	
	1	
R.sp	GSNGWAFGADATANRRGVLLGNPHFPWTTTNRFYQVHLTVPGKLDVMGASIAAFPVVSIGFNKDVWHTVSTGRRFTLF	311
D.rad	GSNGWAFGSEATTNGRGLLGNPHFPWETSNRFYQLHLTLPGQFDVMGASLGMPVNVNIGFNQDVWHTVSTDKRFTLA	284
A.uta	GSNAYGLGAQATVNGSGMVLNPHFPWQGAERFYRMHLKVPGRYDVEGAALIGDPIIEIGHNRTVAWSHTVSTARRFVWH	288
P.aer	GSNAIavgSERSADGKMLLANPHFPWNGAMRFYQMHLLTIPGRLDVMGASLPGLPVNVNIGFSRHLAWHTVDTSSHTFLY	290
	*** . . : : : * : : * . * . * : : * : : * : : * : : * : : * : : * : : * : : *	

```

R.sp      ELKLAEGDPTTYLVDGTPHKMTTTRTVAFDVKLPDGRLERRHTHTFYDTIYGPVLSMPSGGMPWTTQKAYALRDANRNNTRS 391
D.rad     ALTLVPGDPLSYVKDGGQRRRLQRRTAVIEVKTANG-PRLHTRTVYFTPEGPLVNLPAAGLTWTPQYAFALRDANRNNTRM 383
A.uta     RLSLVPGDPTSYVVDGRPERMRARTVTVQTGSG-----PVSRTFHDTRYGPVAVVP-GTFDWTPTATAYAITDVNAGNNRA 383
P.aer     RLALDPKDPRRYLVDGRSLPLEEKSVAIEVRGADGKLSRVEHKVYQSIYGPLVWVP-GKLDWNRSEAYALRDANLENTRV 390
          * *   ** *   **   :   : : : : :   : : : :   ** :   * . : * .   * : * : * *   * : *
          :

R.sp      VDSWLHIGQARDVAGIRQAIG-NLGIPWVNTIATDRNGRALFADVSTTPDVPAEELQRCAPSPLAGKLFKDAGLVLLDGS 470
D.rad     LATWLGFAGAKSVRDIRASLN-VQGIPWVNTIADDRAGSALYADISSSPNVSAQQQACTPPPLA-PLFFAAGLAVLDGS 461
A.uta     FDGWLRMGQAKDVRALKAVLDRHQFLPWVNVIAADARGEALYGDHSVVPRTGALAAACIPAPFQ-PLYASSGQAVLDGS 461
P.aer     LQQWYSINQASDVADLRRRVEALQGIPWVNTLADEQGNALYMNQSVVPYLPKELIPACAIPQLV-----AEGLPALQGG 464
          . * : * . * : : :   : **** : : * * * : : * * :   * . :   * * : * :
          :

R.sp      RGTCNWQVDPASVPVGLVAFARMPVLERDDYVANSNDSSWLTNPAQKLTGFSFVMSVDVPQRLRTRIGLIEIGRRLAGT 550
D.rad     HSACDWKTDPASRVPGLRAPDKMPVLIRQDFVANSNNSAWLANPAAPQTGLDPLVGEVNAQSPRTRMGLEIGRRLSGT 541
A.uta     RSDCALGADPDAAVPGILGPASLPVFRDDYVTNSNDSHWLASPAAPLEGFPRLGNERTPRSLRTRLGLDQIQQLAGT 541
P.aer     DSRCAWSRDPAAAQAGITPAAQLPVLLRRDFVQNSNDSAWLTNPASPLQGFSPLVSQE-KPIGPRARYALSRLQKQP-- 543
          . *   ** :   . :   . : ** * * : * * : * * : * *   * :   : :
          :

R.sp      DGLPGNRIDLPLNQLAMIFSNANLAGQLVLGDLAACKATPAPDAD-----VRDGAALGQWNRTSNADA-RAAHLFREF 630
D.rad     DGLPGRTFDIPTLQATLLRESNLTGEMYAADAALKCS--AGGAE-----LQPACNALAAWDRSSQES-RGAALWREF 619
A.uta     DGLPGKGFTTARLWQVMFGNRMHGAELVRDDLVALCRRQPTATASNGAIVDLTAACIALSRFDERADLDS-RGAHLFTEF 621
P.aer     -----LEAKTLEEMVTANHVFSAQVLPDLLRLCRDN-QGEKS-----LARACAALAQWDRGANLDSGSGFVYFQRF 613
          :   *   :   :   . :   *   * :   .   :   . * ** . : : . : :   . : *
          :

R.sp      WMRAKDIAQVHAVEFDPADPVHTPRGLR-MNDATVRTAVFKALKEAVGAVRKAGFALDAPLGTVQAAHAPDGSIALHGGE 702
D.rad     WRRARAI PNVIYAVPFDPADPVNTPRGLN-TADPAAQTALLGALREAAAALTAAGIPFDAPLGEVQGVVRGGDFISLPGGA 691
A.uta     LAGG-----IRFADTFEVTDPVRTPAEFWNTTDFPRVTALADACNGSPASPSTR-----SVGDIHTDSRGERRIPIHGGR 690
P.aer     MQRFaelDGAWKEPFDAQRPLDTPQGIa-LDREQVATQVRQALADAAAEVEKSGIPD GARWGDQVSTRGQERIAIPGGD 686
          * :   * : ** :   . . * : *   :   .   * : :   * : : **
          :

R.sp      EYEGVLNKLQTLPIGPKGLPVYFG--TSYIQTVTFDDQGFVADAILTYGESTDHASPHAFDQMRAYSGKHWNRLPFSEAA 780
D.rad     EFEGVLNKLIDFNPLAPGGYRGVGNASSYIQTVGFTDSGVQAEAVLTYSQSSNPESPYFSDQTRLFSRSEWVKLPFTQPE 771
A.uta     GEAGTFNVITNPLVPGVGYPQVHVG-TSFMVAVELGPHGSGRQILTYAQSTNPNPWPYADQTVLYSRKGDWDTIKYTEAQ 769
P.aer     GHFGVYNAIQS--VRKGDHLEVVG--TSYIQLVTFPEEGPKARGLLAFSQQSSDPRSPHYRDQTELF SRQQWQTLPFSDRQ 763
          * . : :   :   .   . : : : * :   * .   : : : : : : **   **   : * . *   : : :
          :

R.sp      IAADPALKVMRLSQ--- 794
D.rad     IEADPTRTVVQLSE--- 785
A.uta     IAADPNLRVYRVAQRGR 786
P.aer     IDADPQLQLRSIRE--- 777
          * ***   : :
          :

```

\* = identical residues, : = conserved substitutions, . = semi-conserved substitutions, ! = post-translational processing sites for signal peptide and subunits, - = spacers.

[00029] The coding region of the *gsbA* gene was amplified by PCR. The amplified PCR products were digested, fused in-frame to the coding sequence of the glutathione S-transferase (GST) gene and expressed in *E. coli*. Protein extracts from the recombinant *E. coli* cells were assayed for the ability to inactivate AHL. Protein from *E. coli* expressing GST alone

served as a control. The results demonstrated that GST-QsbA fusion protein effectively eliminated AHL activity. See Figure 1B.

**[00030]** The substrate specificity of QsbA was determined by assaying total soluble protein extracted from the recombinant *E. coli* (pGST-QsbA) for inactivation of AHLs using substrates with acyl chains of differing lengths. QsbA was able to completely inactivate 3-oxo group acyl-HSLs having acyl chains of 8, 10 and 12 carbons. QsbA also strongly inactivated methylene group acyl-HSLs having acyl chains of 8 and 10 carbons. QsbA also inactivated the butyl and hexyl esters of N- $\beta$ -octanoyl-L-homoserine, whereas the AHL-lactonase encoded by *aiiA* was unable to inactivate them. The substrate specificity data indicate that QsbA is an AHL-acylase.

**[00031]** QsbA and *qsbA* provide new tools for down regulation of AHL-mediated biological activities, such as the expression of virulence genes and biofilm differentiation in pathogenic bacteria, both *in vitro* and *in vivo*. The *qsbA* gene, which encodes the AHL inactivation enzyme (QsbA), or a functional fragment, subunit or substantially homologous variant thereof, may be introduced into a plant genome to produce a genetically modified plant with the ability to quench pathogen quorum-sensing signaling. Transgenic plants expressing an enzyme that inactivates AHLs can exhibit a significantly enhanced resistance to infection by bacterial pathogens, even when bacteria are present in high concentrations.

**[00032]** Methods of genetic manipulation and transformation of plant cells are well known in the art, as are methods of regenerating fertile, viable transformed plants. In general, any method of cloning the coding region of *qsbA* or a functional fragment or substantially homologous variant thereof into a suitable expression vector may be used. It is convenient to ligate the *qsbA* coding region into a vector, followed by ligation into a plant transformation vector, however those of skill are well aware of alternative methods to achieve the same results. Any suitable plant transformation vector may be used.

The vector contains the *qsbA* gene, or a functional fragment, subunit or substantially homologous variant thereof, so long as expression of the gene results in a QsbA protein or functional fragment, subunit or variant thereof which inactivates AHL.

[00033] A functional promoter preferably controls expression. Many suitable promoters are known in the art, such that a convenient promoter may easily be selected by a skilled artisan depending on the expression system being used. Such selection of a suitable promoter to achieve the desired level of translational expression is considered routine in the art. For example, it is advantageous to optimize *qsbA* expression by modification of codon usage and coupling to a strong promoter such as the double 35S promoter.

[00034] A suitable marker gene, such as kanamycin resistance, green fluorescent protein or any other convenient marker is advantageously used. Variations of the commonly used and well known methods for transforming plants with a gene, are well within the skill of the ordinary artisan in genetic manipulation of plants. Expression constructs may contain a signal sequence to direct secretion of the expressed QsbA protein, or may lack such a sequence, as desired. The plant transformation vectors containing the *qsbA* gene and a marker gene may be used to transform plant cells using *Agrobacterium*-mediated transformation. *Agrobacterium*-mediated transformation is conveniently used to transform plants with the *qsbA* gene, however any suitable method known in the art may be used, depending on the plant being transformed. For example, certain monocotyledonous plants are more efficiently transformed using other methods such as microprojectile bombardment, vacuum filtration or any other method known in the art to introduce and integrate DNA plasmids or fragments into the plant genome. Those of skill in the art are familiar with means to transform gymnosperms, monocots and dicots. All of these methods known in the art are contemplated for use with this invention.

[00035] After selection for transformants carrying the *qsbA* gene, transgenic plants may be regenerated according to known

methods in the art. Plants selected for a marker gene, for example kanamycin resistance, may be assayed, for example by PCR and DNA gel blot to determine how many copies of the *qsbA* gene are present in the plant tissue. Any suitable method known in the art is contemplated for use with the gene of this invention. *QsbA* enzyme activity may be detected in transgenic plants transformed with the *qsbA* gene by the bioassay method described in Example 2 or by any convenient method.

**[00036]** By "functional fragment, subunit or substantially homologous variant thereof," when referring to a *qsbA* nucleotide sequence, it is meant any fragment, subunit, variant or homologous sequence of *qsbA* (nucleotides 1234-3618 of SEQ ID NO: 1) which encodes a protein or peptide sequence capable of inactivating N-acyl homoserine lactones. "Substantially homologous variants" of a nucleotide sequence generally are those the complement of which hybridizes with *qsbA* under stringent or highly stringent conditions, for example temperatures of about 30°C to about 50°C, for example 30°C, 35°C, 37°C, 40°C, 45°C or 50°C, and/or salt concentrations of about 200 mM to about 1000 mM NaCl or the equivalent ionic strength, for example 200 mM, 250 mM, 300 mM, 400 mM, 500 mM, 750 mM or 1000 mM. The stringency conditions are dependent on the length of the nucleic acid and the base composition of the nucleic acid and can be determined by techniques well known in the art. Those of skill in the art are familiar with these conditions and ranges which are useful. Generally, a substantially homologous nucleotide sequence is at least about 75% homologous to SEQ ID NO: 1 or a fragment or subunit thereof, preferably at least about 85% homologous, and most preferably 90%, 95% or 99% homologous or more.

**[00037]** Those of skill in the art are familiar with the degeneracy of the genetic code, and thus are aware that nucleic acid sequences may be less than 100% homologous and yet encode the same protein or peptide sequence. Such variation in any of the sequences, fragments, subunits or substantially homologous variants also are contemplated as part of this invention.

**[00038]** Peptide and protein sequences which are encompassed by this invention include any sequences encoded by the *qsba* gene, or any fragment, subunit or substantially homologous variant thereof. Such sequences therefore include any functional protein or peptide which retains the ability to inactivate AHL, including protein and peptide fragments of the complete QsbA protein, such as, for example, the sequences of amino acids 36-217 and 233-794 encoding by SEQ ID NO: 1 and substantially homologous variants thereof. A substantially homologous variant of the QsbA protein includes sequences which are at least about 50% homologous, preferably at least about 60% homologous, and most preferably 70%, 80% or 90% homologous or more. Therefore, a protein which is a substantially homologous variant of QsbA is about 50% to about 99.9% homologous with QsbA. Both conservative and non-conservative amino acid substitutions are contemplated, as well as sequences containing non-traditional or modified amino acids such as those known in the art.

**[00039]** The term "fragment" is intended to indicate any portion of the nucleotide of SEQ ID NO: 1 or protein/peptide sequence of SEQ ID NO: 2 which is greater than about 300 nucleotide bases or about 100 amino acids, up to one nucleotide or amino acid less than the entire sequence. The term "subunit" is intended to encompass any functional unit of the QbsA protein, such as, for example, amino acids 36-217 or 233-794 of SEQ ID NO: 2.

**[00040]** A protein or peptide sequence which is considered to inactivate N-acyl homoserine lactones is one which is capable of inactivating at least 55 pmoles N-acyl homoserine lactone (OOHL) per  $\mu$ g protein per minute at 30°C.

**[00041]** It has been previously demonstrated that quenching bacterial quorum sensing by inactivation of N-acyl homoserine lactone with AHL-lactonase stops bacterial infection (9, 10). The gene and protein described here, which is likely an AHL-acylase, represent a new and effective tool for inactivation of AHL signals and thus control bacterial infection. Similarly,

the gene and protein described here targets AHL quorum-sensing signals that regulate expression of pathogenic genes of many bacterial pathogens at a threshold concentration. This tool is applicable to all plant, animal or human diseases where the expression of pathogenic genes of bacterial pathogens is activated by AHL signals, such as, for example, plant pathogens *Erw. carotovora*, *Erw. Chrysanthemi*, *Erw. Stewartii*; human pathogens *P. aeruginosa*, *B. cepacia*; and animal pathogens *X. nematophilus*, *P. fluorescens* (1, 3, 6, 12, 17, 19, 22, 23, 24, 26).



## References

1. Allison, et al., "Extracellular products as mediators of the formation and detachment of *Pseudomonas fluorescens* biofilms," *FEMS Microbiol. Lett.* 167:179-184, 1998.
2. Bassler, et al., "Cross-species induction of luminescence in the quorum-sensing bacterium *Vibrio harveyi*," *J. Bacteriol.* 179:4043-4045, 1997.
3. Beck von Badman, and Ferrand, "Capsular polysaccharide biosynthesis and pathogenicity in *Erwinia stewartii* require induction by an *N*-acyl homoserine lactone autoinducer," *J. Bacteriol.* 177:5000-5008, 1995.
4. Cao and Meighen, "Purification and structural identification of an autoinducer for the luminescence system of *Vibrio harveyi*," *J. Biol. Chem.* 264:21670-21676, 1989.
5. Cha et al., "Production of acyl-homoserine lactone quorum-sensing signals by gram-negative plant associated bacteria," *Mol. Plant Microbe Interact.* 11:1119-1129, 1998.
6. Costa and Loper, "EcbI and EcbR: homologs of LuxI and LuxR affecting antibiotic and exoenzyme production by *Erwinia carotovora* subsp. *betavasculorum*," *Can. J. Microbiol.* 43:1164-1171, 1997.
7. Daumy et al., "Role of protein subunits in *Proteus retigeri* penicillin G acylase," *J. Bacteriol.* 163:1279-1281, 1985.
8. Davies et al., "The involvement of cell-to-cell signals in the development of a bacterial biofilm," *Science* 280:295-298, 1998.
9. Dong et al., "AiiA, an enzyme that inactivates the acyl homoserine lactone quorum-sensing signal and attenuates the virulence of *Erwinia carotovora*," *Proc. Natl. Acad. Sci. USA* 97:3526-3531, 2000.
10. Dong et al., "Quenching quorum sensing-dependent bacterial infection by an *N*-acyl homoserine lactonase," *Nature* 411:813-817, 2001.
11. Dumenyo et al., "Genetic and physiological evidence for the production of *N*-acyl homoserine lactones by *Pseudomonas syringae* pv. *syringae* and other fluorescent plant pathogenic *Pseudomonas* species," *Eur. J. Plant Pathol.* 104:569-582, 1998.
12. Dunphy et al., "A homoserine lactone autoinducer regulates virulence of an insect-pathogenic bacterium, *Xenorhabdus nematophilus* (Enterobacteriaceae)," *J. Bacteriol.* 179:5288-5291, 1997.

13. Eberhard et al., Structural identification of autoinducer of *Photobacterium fischeri* luciferase," *Biochemistry* 20:2444-2449, 1981.
14. Eberl et al., "Involvement of N-acyl-L-homoserine lactone autoinducers in controlling the multicellular behaviour of *Serratia liquefaciens*," *Mol. Microbiol.* 20:127-136, 1996.
15. Fuqua and Winans, "Conserved cis-acting promoter elements are required for density-dependent transcription of *Agrobacterium tumefaciens* conjugal transfer genes," *J. Bacteriol.* 178:435-440, 1996.
16. Inokoshi et al., "Cloning and sequencing of the aculeacin A acylase-encoding gene from *Actinoplanes utahensis* and expression in *Streptomyces lividans*," *Gene* 119:29-35, 1992.
17. Jones et al., "The Lux autoinducer regulates the production of exoenzyme virulence determination in *Erwinia carotovora* and *Pseudomonas aeruginosa*," *EMBO J.* 12:2477-2482, 1993.
18. Leadbetter and Greenberg, "Metabolism of acyl-homoserine lactone quorum sensing signals by *Variovorax paradoxus*," *J. Bacteriol.* 182:6921-6926, 2000.
19. Lewenza et al., "Quorum sensing in *Burkholderia cepacia*: identification of the LuxRI homologs CepRI," *J. Bacteriol.* 181:748-756, 1999.
20. Matsuda and Komatsu, "Molecular cloning and structure of the gene for 7 $\beta$ -(4-carboxybutanamido)cephalosporadic acid acylase from a *Pseudomonas* strain," *J. Bacteriol.* 163:1222-1228, 1985.
21. Matsuda et al., "Nucleotide sequence of the genes for two distinct cephalosporin acylases from a *Pseudomonas* strain," *J. Bacteriol.* 169:5821-5826, 1987.
22. Nasser et al., "Characterization of the *Erwinia chrysanthemi* *expI-expR* locus directing the synthesis of two N-acyl-homoserine lactone signal molecules," *Mol. Microbiol.* 29:1391-1405, 1998.
23. Passador et al., "Expression of *Pseudomonas aeruginosa* virulence genes requires cell-to-cell communication," *Science* 260:1127-1130, 1993.
24. Pearson et al., "Structure of the autoinducer required for expression of *Pseudomonas aeruginosa* virulence genes," *Proc. Natl. Acad. Sci. USA* 91:197-201, 1994
25. Piper et al., "Conjugation factor of *Agrobacterium tumefaciens* regulates Ti plasmid transfer by autoinduction," *Nature* 362:448-450, 1993.

26. Pirhonen et al., "A small diffusible signal molecule is responsible for the global control of virulence and exoenzyme production in the plant pathogen *Erwinia carotovora*," *EMBO J.* 12:2467-2476, 1993.
27. Schumacher et al., "Penicillin acylase from *E. coli*: unique gene-protein relation," *Nucleic Acids Res.* 14:5713-5727, 1986.
28. Staskawicz et al., "Molecular characterization of cloned avirulence genes from race 0 and race 1 of *Pseudomonas syringae* pv. *glycinea*," *J. Bacteriol.* 169:5789-5794, 1987.
29. Takeshima et al., "A deacylation enzyme for aculeacin A, a neutral lipopeptide antibiotic, from *Actinoplanes utahensis*: purification and characterization," *J. Biochem.* 105:606-610, 1989.
30. Verhaert et al., "Molecular cloning and analysis of the gene encoding the thermostable penicillin G acylase from *Alcaligenes faecalis*," *Appl. Env. Microbiol.* 63:3412-3418, 1997.
31. Zhang et al., "Agrobacterium conjugation and gene regulation by *N*-acyl-L-homoserine lactones," *Nature* 362:446-447, 1993.

[00042] The following examples are provided to illustrate the invention described herein and should not be construed to limit the appended claims.

#### EXAMPLES

##### Example 1. Bacterial Isolation.

[00043] A bacterial biofilm sample was collected from a water treatment system and screened to isolate AHL inactivation bacterial strains. The bacterial mixture was suspended in sterilized water with shaking for 1 hour before spreading onto YEB medium (yeast extract, 5 g/l; casein hydrolysate, 10 g/l; NaCl, 5 g/l; sucrose, 5 g/l;  $MgSO_4 \cdot 7H_2O$ , 0.5 g/l and agar, 15 g/l) plates. Individual colonies were restreaked on new plates to ensure purity of the isolates. Bacterial isolates were cultured in LB medium (tryptone, 10 g/L; yeast extract, 5 g/L, and NaCl, 10 g/L; pH7.0) in 1.5-ml Eppendorf™ tubes or 96-well plates at 28°C, with shaking, overnight, and assayed for AHL inactivation activity.

Example 2. AHL Inactivation Bioassay.

[00044] The bacterial culture to be assayed was mixed with an equal volume of fresh medium containing 20  $\mu$ M N- $\beta$ -oxooctanoyl-L-homoserine lactone (OOHL), or another AHL, when specified, to form a reaction mixture. The reaction mixture was incubated at 28°C for 4 to 5 hours, followed by 30 minute sterilization under UV light. Plates containing 20 ml MM agar medium ( $K_2HPO_4$ , 10.5 g/L;  $KH_2PO_4$ , 4.5 g/L;  $MgSO_4 \cdot 7H_2O$ , 0.2 g/L;  $FeSO_4$ , 4.5 mg/L;  $CaCl_2$ , 10 mg/L;  $MnCl_2$ , 2 mg/L;  $(NH_4)_2SO_4$ , 2.0 g/L; mannitol, 2.0 g/L; pH 7.0) supplemented with 5-bromo-4-chloro-3-indolyl  $\beta$ -D-galactopyranoside (X-Gal, 40  $\mu$ g/ml) were prepared. The solidified medium was cut, still inside the plate, into separated slices (approximately 1 cm in width). See Figure 1. Five microliters of sterilized reaction mixture was loaded at the top of an MM agar strip, and then AHL indicator cells (*Agrobacterium tumefaciens* strain NT1 (*traR*; *tra::lacZ749*) (25) 0.7  $\mu$ l cell suspension with an optical density at 600 nm of 0.4) were spotted at progressively further distances from the loaded samples. Plates were incubated at 28°C for 24 hours. A positive result for AHL inactivation is shown by the absence of blue colonies on the slice. A negative result for AHL inactivation is shown by the presence of blue colonies on the slice. For assay of protein for enzyme activity, total soluble bacterial protein was incubated with 20  $\mu$ M of OOHL (or other AHL) at 37°C for 1 hour as the reaction mixture.

Example 3. Identification and Cloning the *qsba* Gene.

[00045] Two bacterial isolates from the biofilm sample with distinct phenotypes, XJ12B and XJ12A, were found to possess the ability to inactivate AHL, with XJ12B showing stronger enzyme activity. The XJ12B isolate was cultured, centrifuged and sonicated. The strongest enzymatic activity was associated with the cell debris fraction rather than the soluble protein and supernatant fractions. These results indicated that the AHL inactivation activity is membrane associated. Sequencing of 16S rRNA was performed to identify the XJ12A and XJ12B isolates.

The 16S rRNA sequences of these isolates showed 97% and 96% identity, respectively, with that of *Ralstonia eutropha*.

**[00046]** To identify the gene encoding for AHL inactivation, a cosmid library of 1600 clones was constructed in *E. coli* with the genomic DNA of *Ralstonia sp.* strain XJ12B. Genomic DNA from the isolated *Ralstonia sp.* strain XJ12B was partially digested with Sau3A. The resulting DNA fragments were ligated to the dephosphorylated BamHI site of cosmid vector pLAFR3 (28). The ligated DNA was packed with Gigapack IIIIXL Packaging Extract (Stratagene) and transfected into *E. coli* DH5alpha. These *E. coli* transfectants were screened for AHL inactivation activity according to the methods described in Example 2 using OOHl as the substrate. Only a single clone (p13H10) was identified as showing AHL inactivation activity (see Figure 1A, slice 1). To subclone the gene encoding the detected activity, cosmid DNA from the positive clone p13H10 was partially digested with Sau3A and fused into BamHI digested cloning vector pGEM-3Zf (+). The plasmids were ligated and transformed into *E. coli*, and the *E. coli* were assayed for the ability to inactivate AHL as described in Example 2. Clone p2B10, which contains a 4 kb insert, had AHL inactivation activity (see Figure 1A, slice 2). The TGS™ Template Generation System F-700 (Finnzymes OY) was used to mutate p2B10 plasmid DNA by randomly inserting the artificial Mu transposon, following the manufacturer's instructions. Plasmid clone p2B10, which contains the 4 kb insert containing the *qsba* gene, was used as a template. Fifteen mutant clones were produced, and none was able to inactivate AHL. Bacterial cultures of *E. coli* DH5α containing pMUG3 and pMUC6 are shown as examples in Figure 1A, slices 3 and 4, respectively. Plasmids were subsequently purified for sequencing using primers supplied in the kit.

Example 4. Sequencing and Sequence analysis of the *qsba* Gene.

**[00047]** Sequencing was performed according to known methods using ABI Prism dRhodamine Terminator Cycle Sequencing Ready Reaction Kit (Perkin-Elmer Applied Biosystems). The 4 kb

fragment from clone p2B10 was completely sequenced and is shown in Table I. The sequence contains an open reading frame of 2385 nucleotides with an ATG start codon and a TGA stop codon (SEQ ID NO: 1, nucleotides 1259-3643). Based on the MU transposon mutagenesis data described in Example 3, this open reading frame is the coding region of the AHL inactivation gene, designated as *qsbA*. A putative ribosome binding site (AGGAGA) is located 6 base pairs upstream of the first ATG start codon (underlined in Table I).

**[00048]** The deduced peptide sequence shows the typical polypeptide primary structure of aculeacin A acylases (AACs) and penicillin G acylases, with signal peptide- $\alpha$  subunit-spacer- $\beta$  subunit organization (16, 30). There are four additional potential start codons located 3, 36, 189 and 384 downstream from the first ATG. The longest open reading frame encodes 794 amino acids, with a predicted molecular weight of 85373 Daltons. The deduced peptide has 78 strongly basic and strong acid amino acid residues and a predicted isoelectric point of 7.48. The first 20 amino acid residues of the assumed open reading frame appear to be a signal peptide.

**[00049]** The peptide sequence of *qsbA* deduced from the open reading frame shares 40-52% identity with AACs from *Deinococcus radiodurans* strain R1, *Actinoplanes utahensis* and a putative acylase from *Pseudomonas aeruginosa*. The AACs' catalyze deacylation of their substrates. These AAC genes are translated as single precursor polypeptide and then processed to the active form of two subunits. By alignment with the peptide sequences from *D. radiodurans* strain R1, *A. utahensis* and *P. aeruginosa*, Table II, the presumed  $\alpha$  and  $\beta$  subunits are located at amino acid positions 36-217 and 233-794, respectively, with a 15 amino acid spacer between them. *QsbA* shares less than 28% homology with penicillin G acylase (20) and cephalosporin acylase (21). See Table II. The amino acid sequence alignment in Table II was analyzed by the Clustal W program available from the European Bioinformatics Institute website (<http://www.ebi.ac.uk>).

Example 5. Expression of the *QsbA* Gene.

[00050] The coding region of the *qsbA* gene was amplified by PCR using a forward primer 5'-CGTGGATCCATGATGCAGGATTCGCCGCTGCGC-3' (SEQ ID NO: 6) and a reverse primer 5'-CGCGAATTCACCGGCAGCCCTCATGCGACAAC-3' (SEQ ID NO: 7) containing BamH1 and EcoR1 restriction sites, respectively. The amplified PCR products were digested using the above restriction enzymes and fused in-frame to the coding sequence of the glutathione S-transferase (GST) gene under the control of the isopropyl  $\beta$ -D-thiogalactopyranoside (IPTG) inducible lac promotor in vector pGEX-2T (Amersham Pharmacia) to generate construct pGST-QsbA. pGST-QsbA was transformed into *E. coli* and expressed.

[00051] Total soluble protein was extracted from the recombinant *E. coli* cells harboring the GST-QsbA-encoding fusion construct according to methods known in the art, based on the methods described in Dong et al. (9), and assayed for AHL inactivation. The total soluble protein from *E. coli* containing GST vector only was used as a control. For the bioassay, 50  $\mu$ l of the soluble protein preparation (20  $\mu$ g/ $\mu$ l) was added to the same volume of 40  $\mu$ M AHL, e.g., OOHL. After a 1 hour incubation at 37°C, the reaction mixture was assayed as described in Example 2. Representative data, shown in Figure 1B, slice 1, indicate that the soluble GST-QsbA fusion protein effectively eliminated OOHL activity.

Example 6. Characterization of the Substrate Spectrum of GST-QsbA Fusion Protein Expressed in *E. coli*.

[00052] To determinate the substrate spectrum of QsbA, total soluble protein extracted from the recombinant *E. coli* (pGST-QsbA) was assayed for inactivation of AHLs with acyl chains of differing lengths according to the methods of Example 2. The following AHLs were synthesized according to known methods as described by Zhang et al. (31): (1) N-octanoyl-L-homoserine lactone (C8HSL, OOHL); (2) N-decanoyl-L-homoserine lactone (C10HSL, DHL); (3) N- $\beta$ -oxohexanoyl-L-homoserine lactone (3-oxo-

C6HSL, OHHL); (4) N- $\beta$ -oxohexanoyl-L-homoserine lactone (3-oxo-C12HSL, OdDHL); (5) N- $\beta$ -oxohexanoyl-L-homoserine lactone (3-oxo-C8HSL, OOHL). The butyl and hexyl esters of N- $\beta$ -oxohexanoyl-L-homoserine were prepared by esterification of N- $\beta$ -oxohexanoyl-L-homoserine lactone with 1-butanol and 1-hexanol respectively, in the presence of small amount of Dowex 50H+ resin (Aldrich). The reaction was conducted at 60°C for 2 hours and the products were purified by silica column chromatography.

**[00053]** QsbA completely inactivated OOHL, N- $\beta$ -oxodecanoyl-L-homoserine (ODHL) and N- $\beta$ -oxododecanoyl-L-homoserine (OdDHL), which have acyl chains of 8, 10 and 12 carbons, respectively, at the concentrations tested (data not shown). QsbA also strongly inactivated N- $\beta$ -octanoyl-L-homoserine(OHL) and N- $\beta$ -decanoyl-L-homoserine(DHL), which have acyl chains of 8 and 10 carbons, respectively (data not shown). However, under the same reaction conditions, QsbA had less inactivating activity for N- $\beta$ -oxohexanoyl-L-homoserine (OHHL), which has an acyl chain of 6 carbons (data not shown). The total soluble protein extract from control *E.coli* (pGST) did not show any activity against AHLs (data not shown).

**[00054]** QsbA also completely inactivated the butyl and hexyl esters of N- $\beta$ -octanoyl-L-homoserine (data not shown). These two esters of N- $\beta$ -octanoyl-L-homoserine showed comparable induction activity with OOHL when assayed with the AHL reporter strain *A. tumefaciens* NT1 (*traR*; *tra::lacZ749*) (data not shown). AHL-lactonase (encoded by *aiiA*) did not inactivate these substrates. These substrate specificity data are consistent with identification of QsbA as an AHL-acylase.

**Example 7.** Purification of AHL-acylase encoded by the *qsbA* gene

**[00055]** The GST-[AHL-acylase] fusion protein was purified using a glutathione Sepharose 4B affinity column following the manufacturer's instructions (Pharmacia). AHL-acylase was cleaved by digestion with thrombin (Sigma). Protein concentration was determined by measuring OD<sub>280</sub>.



[00056] The purified AHL-acylase was incubated with OOHL for 20 minutes and the relative enzyme activity was measured by determining the residual OOHL in the reaction mixture, which contained 8  $\mu$ M OOHL and about 0.6  $\mu$ g AHL-acylase in a total reaction volume of 50 ml 1x PBS buffer. The reactions were stopped by addition of 1% SDS before loading on the assay plate. Determination of the OOHL activity was carried out in quadruplicate. AHL-acylase degraded OOHL in a range of temperatures from 22-42°C at pH 7.0. See Figure 2. The optimal temperature for enzyme activity was found to be approximately 28°C. Reaction temperature higher than 42°C decreased enzyme activity sharply. The optimal pH for enzyme activity also was determined. The AHL-acylase has a relatively narrow optimal pH range from pH 6.5 to 7.5. See Figure 2. The time course of OOHL inactivation by the purified AHL-acylase was determined at 30°C. See Figure 3. After 10 minutes, more than 82% OOHL had been degraded; the reaction rate was estimated to be about 55 pmols per  $\mu$ g AHL-acylase per minute.

## CLAIMS:

1. A composition of matter which comprises a nucleic acid according to SEQ ID NO: 1.
2. A composition of matter which comprises a nucleic acid selected from the group consisting of nucleotides 1234-3618 of SEQ ID NO: 1, a fragment thereof and a substantially homologous variant thereof.
3. A nucleic acid according to claim 2 which comprises nucleotides 1234-3618 of SEQ ID NO: 1.
4. A composition of matter which comprises a peptidic sequence selected from the group consisting of a peptidic sequence according to SEQ ID NO: 2, a fragment thereof and a substantially homologous variant thereof.
5. A composition of matter which comprises a peptidic sequence encoded by a nucleic acid selected from the group consisting of nucleotides 1234-3618 of SEQ ID NO: 1, a fragment thereof and a substantially homologous variant thereof.
6. A composition of matter which comprises a peptidic sequence selected from the group consisting of SEQ ID NO: 2, a fragment thereof, a subunit thereof and a substantially homologous variant thereof.
7. A composition of matter according to claim 6 which comprises a peptidic sequence according to SEQ ID NO: 2.
8. A composition of matter according to claim 6 which comprises a peptidic sequence comprising amino acids 36-217 of SEQ ID NO: 2.

9. A composition of matter according to claim 6 which comprises a peptidic sequence comprising amino acids 233-794 of SEQ ID NO: 2.

10. A composition of matter according to claim 6 which inactivates AHL.

11. A method of modulating AHL signaling activity which comprises contacting said AHL with a composition of matter according to any one of claims 5-10.

12. A transgenic plant harboring a nucleic acid selected from the group consisting of nucleotides 1234-3618 of SEQ ID NO: 1, a fragment thereof and a substantially homologous variant thereof.

13. A transgenic non-human animal harboring a nucleic acid selected from the group consisting of nucleotides 1234-3618 of SEQ ID NO: 1, a fragment thereof and a substantially homologous variant thereof.

14. A method of controlling a bacterial disease in a mammal in need thereof which comprises administering to said mammal a composition of matter according to any one of claims 5-10, wherein the expression of pathogenic genes of said bacteria are regulated by AHL signals.

15. A method of claim 14 wherein said mammal is a human.

16. A method of controlling a bacterial disease in a plant in need thereof which comprises administering to said plant a composition of matter according to any one of claims 5-10, wherein the expression of pathogenic genes of said bacteria are regulated by AHL signals.

17. A method of controlling a bacterial disease in a mammal in need thereof which comprises administering to said mammal a

composition of matter of claim 2 and its peptide product, wherein the expression of pathogenic genes of said bacteria are regulated by AHL signals.

18. A method of claim 17 wherein said mammal is a human.

19. A method of controlling a bacterial disease in a plant in need thereof which comprises administering to said plant a composition of matter of claim 2, wherein the expression of pathogenic genes of said bacteria are regulated by AHL signals.

20. A method of controlling a bacterial disease in a plant using any bacterial species containing the composition of matter of claim 2.

## RALSTONIA AHL-ACYLASE GENE

## ABSTRACT OF THE DISCLOSURE

[00058] This invention provides a gene, *qsba*, which encodes a protein useful for inactivating certain bacterial quorum-sensing signal molecules (N-acyl homoserine lactones) which participate in bacterial virulence and biofilm differentiation pathways. This gene was isolated from *Ralstonia* sp., strain XJ12B. The invention also provides the QsbA protein, which possesses N-acyl homoserine lactone inactivating activity.

2577-154.PCT

Fig. 1

1/2




FIG. 1A




FIG. 1B




FIG. 2A




FIG. 2B




FIG. 3

## SEQUENCE LISTING

<110> Zhang, Lian Hui

Lin, Yi Han

Xu, Jin Ling

<120> RALSTONIA AHL-ACYLASE GENE

<130> 2577-154 PCT

<160> 7

<170> PatentIn version 3.0

<210> 1

<211> 3743

<212> DNA

<213> Ralstonia sp.

<220>

<221> N\_region

<222> (16)..(16)

<223> unknown

<400> 1  
gtttgggaaa gtgggnagcg cgctgtgcag cgccccgcc ctcagccgcg cagctcggcg 60  
cgcaccgaat gcgcgcgccg gtgggcgccc ggcggtggc cggtgtggcg ccggatcagg 120  
cgccggaagg cggacatgtc gtgataaccg cactgttcgg cgattgccgt caggctcagc 180  
gtgctgactt ccagcaggtg gcaggcgcg cccacgcgca gccggtgcag caattgcagc 240



ggcgaggtgc ccaggggtctt ggtgaaatgc cgcagcagcg tgcgctcgct ggtcgaggcg 300  
 gcggcgggcca gcttggccag gtcgaacggc tcgtgcaggt gctgctgcag gtagcgccgc 360  
 gcccgagta ccacgctggt gcgcatggcg ggcttgctgc gcagccagat ggcggtggac 420  
 tcaccgcgcg acgggtggtc gagcacggcc tggccgaggg tgcgtgccag ccgggtgtcg 480  
 gccaggcggc cgaccaggcg ctgctgagc gccacgccgt gctccatcgc gcgcgccgtc 540  
 agcacgttgc cgctgctgac gatggcctgc tccgccacca ccttcagctg cgggtagttg 600  
 ccgtgcagcc agccggcgat cagccacgtc accgtcaagc gccggccggc gggcagcgcg 660  
 ccggccagca gcgccacgcc ggtgaaggac gagggcacca agcctgccgg cgtccaggta 720  
 gcgcgggatg gtggcgcggt cccactccag cagggccagg cgctgctcca gcgtgctgat 780  
 gtggtcgaaa tgcaggggcg ggacgaccag cgcgtcgccc agcgcggcgt cggccgcccg 840  
 cagcggttg cagcggcagg ccaggggtctc ggcgggcgcc tgccagcggg ccgggtcgcg 900  
 cgcgaccagc cgccaccga acaccgggtt ggcggcacgc gcacgcttgc cggcatgcat 960  
 ggaggcgagc gcattggcca cgccgaggggt gtcggcgacg gtcgccagcg tggagaggcc 1020  
 ggcgtcggga aaggtcagca ggtcgatgtc ggcacccga aagtataggg gaggcgggcg 1080  
 gaggcctcct gcgtggcggg attgaccca actctggcg gaataacctt ttctccggg 1140  
 cgggccccag tcgacgatac ggcggtggct gcgcctgcgc gccgccgcaa gactagagcg 1200  
 acacaagaca agaccgacaa caggagacaa cgcatgatgc agggattcgc gctgcgcggc 1260  
 acgctcgcca tggccgcgct cgcggcgctg gccggctgcg ccagttccac cgatggccgc 1320  
 tgggggtcgc tcagcgacac cggcctgtcc gccgagatcc gccgcaccgg cttcggcatt 1380  
 ccgcacatcc gcgccaacga ctacgccagc ctcggtatg gcatggccta tgcctacgcg 1440  
 caggacaacc tgtgctgct ggccgaccag gtggtcaccg tcaacggcga gcgctcgaag 1500  
 accttcgggc ccgagggcac cgtgacggtc tcgttcaagc cgatcccaa cctgcagtcg 1560  
 gacgccttct tcaagggcat cttcgacgag gacggcctgc gcgccggtta tgcgcagatg 1620  
 tcgcccagg cgcgcgagct gctgcgcggc tacatcgccg gcttcaaccg ctatctcaag 1680  
 gacacgccgc ccgccaactt cccggccgcc tgccgcaatg ccgcctgggt gcgtccgctc 1740  
 acgctgggcg acatgatgcg catggggcga gagaaggcga tccaggccag cgccggcgcc 1800  
 atgctggcg gcacgtcgc cgcgagccg ccgggcccga cgccggtggc cgagcgcgag 1860  
 attccgccgc aggcgctga caccgtggcg ctggaccgcg aactgcagct gcgcgacatg 1920  
 ccgatcggct ccaacggctg ggccttcggc gctgacgcca ccgccaaccg gcgcggcggtg 1980

ctgctcggca atccgcactt cccgtggacg accaccaacc gcttctacca ggtccacctg	2040
acgggtgcccg gcaagctcga cgtgatgggc gcctcgatcg cggccttccc ggtgggtgagc	2100
atcggcttca acaaggacgt ggcgtggacg cacaccgtct ccaccggccg ccgcttcacc	2160
ttgttcgaac tgaagctggc cgaaggcgac ccgaccacct acctggtcga cggcacgccg	2220
cacaagatga ccaccgcac ggtcgccttc gacgtcaagc tgccggacgg ccgcctcgag	2280
cgccgcacgc acaccttcta cgacaccatc tacggcccgg tgctgtcgat gccgagcggc	2340
ggcatgccgt ggaccacgca gaaggcctac gccctgcgcg acgccaaccg caacaacacg	2400
cgctcggtcg acagctggct gcatatcggg caggcccggg acgtggccgg catccgccag	2460
gccatcggca acctgggcat tccctgggtc aacaccatcg ccaccgaccg caacggccgc	2520
gcgctgttcg ccgacgtgtc gaccacgccg gacgtgccgg ccgcggagct ccagcgctgt	2580
gccccgtcgc cgctggccgg caaactcttc aaggacgcgg gcctgggtgct gctcgacggc	2640
tcgcgcggca cctgcaactg gcaggtcgat ccggcttcgc cggtaaccgg gctgggtggc	2700
cccgcgcgca tgccggtgct cgagcgcgac gactacgtcg ccaacagcaa tgacagctcc	2760
tggtcgacca accccgcgca aaagctgacc ggcttctcgc cggatgatgg ctcggtcgac	2820
gtaccgcagc ggctgcgcac gcgcatcggc ctgatcgaga tcggccgccc cctggccggc	2880
accgacggac tgcccggcaa ccgcatcgat ctgccgaacc tgcaggcgat gatcttcagc	2940
aatgccaaacc tggcgggaca actgggtgctg ggcgacctgc tcgcggcatg caaggccacg	3000
ccggccccgg atgccgacgt gcgcgacggc tgccgcgccc tcggccagtg gaaccgcacc	3060
agcaacgccg acgcccgcgc cgcgcacctg ttccgcgagt tctggatgag cgccaaggac	3120
atcgcgcagg tgcacgccgt cgagttcgac ccggccgacc cggccacac gccgcgcggc	3180
ctgcgcatga acgacgcgac ggtacgcacg gcggtgttca aggcgctgaa ggaagccgtg	3240
ggcgcgggtg gcaaggcggg cttcgcgctg gatgcgccgc tgggcacggc acaggccgcg	3300
cacgcaccgg acggctccat cgccctgcac ggccggcgagg aatacgaagg cgtgctcaac	3360
aagctgcaaa ccctgccgat cgggccgaag gggctgccgg tgtatttcgg caccagctac	3420
atccagaccg tgaccttcga cgaccagggc ccggtcgccg acgccatcct cacctacggc	3480
gaatcgaccg accacgcctc gccgcacgcg ttcgaccaga tgcgtgcgta ctcgggcaag	3540
cactggaacc ggctgccgtt ctccgaagcg gccatcgccg ccgatccggc gctgaagggtg	3600
atgcggttgt cgcagtgagg gctgccggtg cctggaaaaa cgccccgctt gtgcggggcg	3660
tttttttgcc agtgtgaatg gctcaatcgt gttggaaacc gcatccggac atgactgtat	3720

3743

<213> Ralstonia sp.

Glu Ile Pro Pro Gln Ala Val Asp Thr Val Ala Leu Asp Arg Glu Leu  
210 215 220

Gln Leu Arg Asp Met Pro Ile Gly Ser Asn Gly Trp Ala Phe Gly Ala  
 225 230 235 240  
 Asp Ala Thr Ala Asn Arg Arg Gly Val Leu Leu Gly Asn Pro His Phe  
 245 250 255  
 Pro Trp Thr Thr Thr Asn Arg Phe Tyr Gln Val His Leu Thr Val Pro  
 260 265 270  
 Gly Lys Leu Asp Val Met Gly Ala Ser Ile Ala Ala Phe Pro Val Val  
 275 280 285  
 Ser Ile Gly Phe Asn Lys Asp Val Ala Trp Thr His Thr Val Ser Thr  
 290 295 300  
 Gly Arg Arg Phe Thr Leu Phe Glu Leu Lys Leu Ala Glu Gly Asp Pro  
 305 310 315 320  
 Thr Thr Tyr Leu Val Asp Gly Thr Pro His Lys Met Thr Thr Arg Thr  
 325 330 335  
 Val Ala Phe Asp Val Lys Leu Pro Asp Gly Arg Leu Glu Arg Arg Thr  
 340 345 350  
 His Thr Phe Tyr Asp Thr Ile Tyr Gly Pro Val Leu Ser Met Pro Ser  
 355 360 365  
 Gly Gly Met Pro Trp Thr Thr Gln Lys Ala Tyr Ala Leu Arg Asp Ala  
 370 375 380  
 Asn Arg Asn Asn Thr Arg Ser Val Asp Ser Trp Leu His Ile Gly Gln  
 385 390 395 400  
 Ala Arg Asp Val Ala Gly Ile Arg Gln Ala Ile Gly Asn Leu Gly Ile  
 405 410 415  
 Pro Trp Val Asn Thr Ile Ala Thr Asp Arg Asn Gly Arg Ala Leu Phe  
 420 425 430  
 Ala Asp Val Ser Thr Thr Pro Asp Val Pro Ala Ala Glu Leu Gln Arg  
 435 440 445  
 Cys Ala Pro Ser Pro Leu Ala Gly Lys Leu Phe Lys Asp Ala Gly Leu  
 450 455 460  
 Val Leu Leu Asp Gly Ser Arg Gly Thr Cys Asn Trp Gln Val Asp Pro  
 465 470 475 480  
 Ala Ser Pro Val Pro Gly Leu Val Ala Pro Ala Arg Met Pro Val Leu  
 485 490 495  
 Glu Arg Asp Asp Tyr Val Ala Asn Ser Asn Asp Ser Ser Trp Leu Thr  
 500 505 510  
 Asn Pro Ala Gln Lys Leu Thr Gly Phe Ser Pro Val Met Gly Ser Val  
 515 520 525

Asp Val Pro Gln Arg Leu Arg Thr Arg Ile Gly Leu Ile Glu Ile Gly  
 530 535 540  
 Arg Arg Leu Ala Gly Thr Asp Gly Leu Pro Gly Asn Arg Ile Asp Leu  
 545 550 555 560  
 Pro Asn Leu Gln Ala Met Ile Phe Ser Asn Ala Asn Leu Ala Gly Gln  
 565 570 575  
 Leu Val Leu Gly Asp Leu Leu Ala Ala Cys Lys Ala Thr Pro Ala Pro  
 580 585 590  
 Asp Ala Asp Val Arg Asp Gly Cys Ala Ala Leu Gly Gln Trp Asn Arg  
 595 600 605  
 Thr Ser Asn Ala Asp Ala Arg Ala Ala His Leu Phe Arg Glu Phe Trp  
 610 615 620  
 Met Arg Ala Lys Asp Ile Ala Gln Val His Ala Val Glu Phe Asp Pro  
 625 630 635 640  
 Ala Asp Pro Val His Thr Pro Arg Gly Leu Arg Met Asn Asp Ala Thr  
 645 650 655  
 Val Arg Thr Ala Val Phe Lys Ala Leu Lys Glu Ala Val Gly Ala Val  
 660 665 670  
 Arg Lys Ala Gly Phe Ala Leu Asp Ala Pro Leu Gly Thr Val Gln Ala  
 675 680 685  
 Ala His Ala Pro Asp Gly Ser Ile Ala Leu His Gly Gly Glu Glu Tyr  
 690 695 700  
 Glu Gly Val Leu Asn Lys Leu Gln Thr Leu Pro Ile Gly Pro Lys Gly  
 705 710 715 720  
 Leu Pro Val Tyr Phe Gly Thr Ser Tyr Ile Gln Thr Val Thr Phe Asp  
 725 730 735  
 Asp Gln Gly Pro Val Ala Asp Ala Ile Leu Thr Tyr Gly Glu Ser Thr  
 740 745 750  
 Asp His Ala Ser Pro His Ala Phe Asp Gln Met Arg Ala Tyr Ser Gly  
 755 760 765  
 Lys His Trp Asn Arg Leu Pro Phe Ser Glu Ala Ala Ile Ala Ala Asp  
 770 775 780  
 Pro Ala Leu Lys Val Met Arg Leu Ser Gln  
 785 790

&lt;210&gt; 3

&lt;211&gt; 785

&lt;212&gt; PRT

&lt;213&gt; D. radiodurans

&lt;400&gt; 3

Met	Ser	Arg	Ser	Pro	Phe	Ser	Ser	Val	Ser	Leu	Pro	Ala	Arg	Leu	Leu	1	5	10	15
Leu	Gly	Ser	Leu	Leu	Leu	Gly	Pro	Leu	Met	Leu	Gly	Gly	Ala	Ala	Ser	20	25	30	
Ala	Gln	Thr	Tyr	Gln	Val	Gln	Ile	Gln	Arg	Thr	Ala	His	Gly	Ile	Pro	35	40	45	
His	Ile	Gln	Ala	Ser	Asp	Leu	Gly	Gly	Ile	Gly	Tyr	Gly	Val	Gly	Tyr	50	55	60	
Ser	Tyr	Ala	Gln	Asp	Asn	Leu	Cys	Leu	Leu	Ala	Asp	Gln	Val	Met	Thr	65	70	75	80
Val	Arg	Gly	Glu	Arg	Ser	Lys	Phe	Leu	Gly	Ala	Glu	Gly	Lys	Thr	Val	85	90	95	
Val	Gly	Phe	Gln	Pro	Val	Asn	Asn	Leu	Asp	Ser	Asp	Val	Phe	Phe	Lys	100	105	110	
Thr	Val	Ile	Glu	Pro	Gly	Arg	Leu	Gln	Ala	Gly	Tyr	Arg	Asp	Gln	Pro	115	120	125	
Gln	Ile	Leu	Ala	Leu	Met	Arg	Gly	Tyr	Val	Ala	Gly	Val	Asn	Arg	Tyr	130	135	140	
Leu	Arg	Asp	Thr	Pro	Pro	Glu	Gln	Trp	Pro	Ser	Ala	Cys	Arg	Asn	Ala	145	150	155	160
Asp	Trp	Val	Arg	Pro	Leu	Thr	Glu	Leu	Asp	Val	Met	Arg	Leu	Gly	Glu	165	170	175	
Glu	Lys	Ala	Ile	Gln	Ala	Ser	Ala	Gly	Ala	Met	Val	Ser	Ala	Ile	Thr	180	185	190	
Ser	Ala	Arg	Pro	Pro	Gln	Ala	Gly	Ala	Ser	Thr	Ala	Ala	Pro	Arg	Pro	195	200	205	
Asp	Leu	Ala	Ala	Phe	Asn	Arg	Gln	Tyr	Arg	Phe	Asn	Asp	Leu	Pro	Ile	210	215	220	
Gly	Ser	Asn	Gly	Trp	Ala	Phe	Gly	Ser	Glu	Ala	Thr	Thr	Asn	Gly	Arg	225	230	235	240

Gly	Leu	Leu	Leu	Gly	Asn	Pro	His	Phe	Pro	Trp	Glu	Thr	Ser	Asn	Arg	245	250	255
Phe	Tyr	Gln	Leu	His	Leu	Thr	Leu	Pro	Gly	Gln	Phe	Asp	Val	Met	Gly	260	265	270
Ala	Ser	Leu	Gly	Gly	Met	Pro	Val	Val	Asn	Ile	Gly	Phe	Asn	Gln	Asp	275	280	285
Val	Ala	Trp	Thr	His	Thr	Val	Ser	Thr	Asp	Lys	Arg	Phe	Thr	Leu	Ala	290	295	300
Ala	Leu	Thr	Leu	Val	Pro	Gly	Asp	Pro	Leu	Ser	Tyr	Val	Lys	Asp	Gly	305	310	315
Gln	Gln	Arg	Arg	Leu	Gln	Arg	Arg	Thr	Ala	Val	Ile	Glu	Val	Lys	Thr	325	330	335
Ala	Asn	Gly	Pro	Arg	Leu	His	Thr	Arg	Thr	Val	Tyr	Phe	Thr	Pro	Glu	340	345	350
Gly	Pro	Leu	Val	Asn	Leu	Pro	Ala	Ala	Gly	Leu	Thr	Trp	Thr	Pro	Gln	355	360	365
Tyr	Ala	Phe	Ala	Leu	Arg	Asp	Ala	Asn	Arg	Asn	Asn	Thr	Arg	Met	Leu	370	375	380
Ala	Thr	Trp	Leu	Gly	Phe	Ala	Gly	Ala	Lys	Ser	Val	Arg	Asp	Ile	Arg	385	390	395
Ala	Ser	Leu	Asn	Val	Gln	Gly	Ile	Pro	Trp	Val	Asn	Thr	Ile	Ala	Ala	405	410	415
Asp	Arg	Ala	Gly	Ser	Ala	Leu	Tyr	Ala	Asp	Ile	Ser	Ser	Ser	Pro	Asn	420	425	430
Val	Ser	Ala	Ala	Gln	Gln	Gln	Ala	Cys	Thr	Pro	Pro	Pro	Leu	Ala	Pro	435	440	445
Leu	Phe	Pro	Ala	Ala	Gly	Leu	Ala	Val	Leu	Asp	Gly	Ser	His	Ser	Ala	450	455	460
Cys	Asp	Trp	Lys	Thr	Asp	Pro	Ala	Ser	Arg	Val	Pro	Gly	Leu	Arg	Ala	465	470	475
Pro	Asp	Lys	Met	Pro	Val	Leu	Ile	Arg	Gln	Asp	Phe	Val	Ala	Asn	Ser	485	490	495
Asn	Asn	Ser	Ala	Trp	Leu	Ala	Asn	Pro	Ala	Ala	Pro	Gln	Thr	Gly	Leu	500	505	510
Asp	Pro	Leu	Val	Gly	Glu	Val	Asn	Ala	Pro	Gln	Ser	Pro	Arg	Thr	Arg	515	520	525
Met	Gly	Leu	Leu	Glu	Ile	Gly	Arg	Arg	Leu	Ser	Gly	Thr	Asp	Gly	Leu	530	535	540

Pro Gly Arg Thr Phe Asp Ile Pro Thr Leu Gln Ala Thr Leu Leu Arg  
 545 550 555 560  
 Glu Ser Asn Leu Thr Gly Glu Met Tyr Ala Ala Asp Ala Ala Lys Leu  
 565 570 575  
 Cys Gln Ser Ala Gly Gly Ala Glu Leu Gln Pro Ala Cys Asn Ala Leu  
 580 585 590  
 Ala Ala Trp Asp Arg Arg Ser Ser Gln Glu Ser Arg Gly Ala Ala Leu  
 595 600 605  
 Trp Arg Glu Phe Trp Arg Arg Ala Arg Ala Ile Pro Asn Val Tyr Ala  
 610 615 620  
 Val Pro Phe Asp Pro Ala Asp Pro Val Asn Thr Pro Arg Gly Leu Asn  
 625 630 635 640  
 Thr Ala Asp Pro Ala Ala Gln Thr Ala Leu Leu Gly Ala Leu Arg Glu  
 645 650 655  
 Ala Ala Ala Ala Leu Thr Ala Ala Gly Ile Pro Phe Asp Ala Pro Leu  
 660 665 670  
 Gly Glu Val Gln Gly Val Val Arg Gly Gly Asp Phe Ile Ser Leu Pro  
 675 680 685  
 Gly Gly Ala Glu Phe Glu Gly Val Leu Asp Lys Ile Asp Phe Asn Pro  
 690 695 700  
 Leu Ala Pro Gly Gly Tyr Arg Gly Val Val Gly Asn Ala Ser Ser Tyr  
 705 710 715 720  
 Ile Gln Thr Val Gly Phe Thr Asp Ser Gly Val Gln Ala Glu Ala Val  
 725 730 735  
 Leu Thr Tyr Ser Gln Ser Ser Asn Pro Glu Ser Pro Tyr Phe Ser Asp  
 740 745 750  
 Gln Thr Arg Leu Phe Ser Arg Ser Glu Trp Val Lys Leu Pro Phe Thr  
 755 760 765  
 Gln Pro Glu Ile Glu Ala Asp Pro Thr Arg Thr Val Val Gln Leu Ser  
 770 775 780  
 Glu  
 785



&lt;210&gt; 4

&lt;211&gt; 786

&lt;212&gt; PRT

&lt;213&gt; A. utahensis

&lt;400&gt; 4

```

Met Thr Ser Ser Tyr Met Arg Leu Lys Ala Ala Ala Ile Ala Phe Gly
1          5          10          15
Val Ile Val Ala Thr Ala Ala Val Pro Ser Pro Ala Ser Gly Arg Glu
20          25          30
His Asp Gly Gly Tyr Ala Ala Leu Ile Arg Arg Ala Ser Tyr Gly Val
35          40          45
Pro His Ile Thr Ala Asp Asp Phe Gly Ser Leu Gly Phe Gly Val Gly
50          55          60
Tyr Val Gln Ala Glu Asp Asn Ile Cys Val Ile Ala Glu Ser Val Val
65          70          75          80
Thr Ala Asn Gly Glu Arg Ser Arg Trp Phe Gly Ala Thr Gly Pro Asp
85          90          95
Asp Ala Asp Val Arg Thr Thr Ser Ser Thr Gln Ala Ile Asp Asp Arg
100         105         110
Val Ala Glu Arg Leu Leu Glu Gly Pro Arg Asp Gly Val Arg Ala Pro
115         120         125
Cys Asp Asp Val Arg Asp Gln Met Arg Gly Phe Val Ala Gly Tyr Asn
130         135         140
His Phe Leu Arg Arg Thr Gly Val His Arg Leu Thr Asp Pro Ala Cys
145         150         155         160
Arg Gly Lys Ala Trp Val Arg Pro Leu Ser Glu Ile Asp Leu Trp Arg
165         170         175
Thr Ser Trp Asp Ser Met Val Arg Ala Gly Ser Gly Ala Leu Leu Asp
180         185         190
Gly Ile Val Ala Ala Thr Pro Pro Thr Ala Ala Gly Pro Ala Ser Ala
195         200         205
Pro Glu Ala Pro Asp Ala Ala Ala Ile Ala Ala Ala Leu Asp Gly Thr
210         215         220
Ser Ala Gly Ile Gly Ser Asn Ala Tyr Gly Leu Gly Ala Gln Ala Thr
225         230         235         240

```

Val	Asn	Gly	Ser	Gly	Met	Val	Leu	Ala	Asn	Pro	His	Phe	Pro	Trp	Gln
				245					250					255	
Gly	Ala	Glu	Arg	Phe	Tyr	Arg	Met	His	Leu	Lys	Val	Pro	Gly	Arg	Tyr
			260					265					270		
Asp	Val	Glu	Gly	Ala	Ala	Leu	Ile	Gly	Asp	Pro	Ile	Ile	Glu	Ile	Gly
		275					280					285			
His	Asn	Arg	Thr	Val	Ala	Trp	Ser	His	Thr	Val	Ser	Thr	Ala	Arg	Arg
	290					295					300				
Phe	Val	Trp	His	Arg	Leu	Ser	Leu	Val	Pro	Gly	Asp	Pro	Thr	Ser	Tyr
305					310					315					320
Tyr	Val	Asp	Gly	Arg	Pro	Glu	Arg	Met	Arg	Ala	Arg	Thr	Val	Thr	Val
				325					330					335	
Gln	Thr	Gly	Ser	Gly	Pro	Val	Ser	Arg	Thr	Phe	His	Asp	Thr	Arg	Tyr
			340					345					350		
Gly	Pro	Val	Ala	Val	Val	Pro	Gly	Thr	Phe	Asp	Trp	Thr	Pro	Ala	Thr
		355					360					365			
Ala	Tyr	Ala	Ile	Thr	Asp	Val	Asn	Ala	Gly	Asn	Asn	Arg	Ala	Phe	Asp
	370					375					380				
Gly	Trp	Leu	Arg	Met	Gly	Gln	Ala	Lys	Asp	Val	Arg	Ala	Leu	Lys	Ala
385					390					395					400
Val	Leu	Asp	Arg	His	Gln	Phe	Leu	Pro	Trp	Val	Asn	Val	Ile	Ala	Ala
				405					410					415	
Asp	Ala	Arg	Gly	Glu	Ala	Leu	Tyr	Gly	Asp	His	Ser	Val	Val	Pro	Arg
			420					425					430		
Val	Thr	Gly	Ala	Leu	Ala	Ala	Ala	Cys	Ile	Pro	Ala	Pro	Phe	Gln	Pro
		435					440					445			
Leu	Tyr	Ala	Ser	Ser	Gly	Gln	Ala	Val	Leu	Asp	Gly	Ser	Arg	Ser	Asp
	450					455					460				
Cys	Ala	Leu	Gly	Ala	Asp	Pro	Asp	Ala	Ala	Val	Pro	Gly	Ile	Leu	Gly
465					470					475					480
Pro	Ala	Ser	Leu	Pro	Val	Arg	Phe	Arg	Asp	Asp	Tyr	Val	Thr	Asn	Ser
				485					490					495	
Asn	Asp	Ser	His	Trp	Leu	Ala	Ser	Pro	Ala	Ala	Pro	Leu	Glu	Gly	Phe
			500					505					510		
Pro	Arg	Ile	Leu	Gly	Asn	Glu	Arg	Thr	Pro	Arg	Ser	Leu	Arg	Thr	Arg
		515					520					525			
Leu	Gly	Leu	Asp	Gln	Ile	Gln	Gln	Arg	Leu	Ala	Gly	Thr	Asp	Gly	Leu
	530					535					540				

Pro Gly Lys Gly Phe Thr Thr Ala Arg Leu Trp Gln Val Met Phe Gly  
 545 550 555 560  
 Asn Arg Met His Gly Ala Glu Leu Val Arg Asp Asp Leu Val Ala Leu  
 565 570 575  
 Cys Arg Arg Gln Pro Thr Ala Thr Ala Ser Asn Gly Ala Ile Val Asp  
 580 585 590  
 Leu Thr Ala Ala Cys Thr Ala Leu Ser Arg Phe Asp Glu Arg Ala Asp  
 595 600 605  
 Leu Asp Ser Arg Gly Ala His Leu Phe Thr Glu Phe Leu Ala Gly Gly  
 610 615 620  
 Ile Arg Phe Ala Asp Thr Phe Glu Val Thr Asp Pro Val Arg Thr Pro  
 625 630 635 640  
 Ala Pro Phe Trp Asn Thr Thr Asp Pro Arg Val Arg Thr Ala Leu Ala  
 645 650 655  
 Asp Ala Cys Asn Gly Ser Pro Ala Ser Pro Ser Thr Arg Ser Val Gly  
 660 665 670  
 Asp Ile His Thr Asp Ser Arg Gly Glu Arg Arg Ile Pro Ile His Gly  
 675 680 685  
 Gly Arg Gly Glu Ala Gly Thr Phe Asn Val Ile Thr Asn Pro Leu Val  
 690 695 700  
 Pro Gly Val Gly Tyr Pro Gln Val Val His Gly Thr Ser Phe Val Met  
 705 710 715 720  
 Ala Val Glu Leu Gly Pro His Gly Pro Ser Gly Arg Gln Ile Leu Thr  
 725 730 735  
 Tyr Ala Gln Ser Thr Asn Pro Asn Ser Pro Trp Tyr Ala Asp Gln Thr  
 740 745 750  
 Val Leu Tyr Ser Arg Lys Gly Trp Asp Thr Ile Lys Tyr Thr Glu Ala  
 755 760 765  
 Gln Ile Ala Ala Asp Pro Asn Leu Arg Val Tyr Arg Val Ala Gln Arg  
 770 775 780  
 Gly Arg  
 785

&lt;210&gt; 5

&lt;211&gt; 777

&lt;212&gt; PRT

&lt;213&gt; P. aeruginosa

&lt;400&gt; 5

```

Met Ser Arg Pro Phe Arg Pro Pro Leu Cys Arg Glu Thr Thr Ser Met
1          5          10          15

Gly Met Arg Thr Val Leu Thr Gly Leu Ala Gly Met Leu Leu Gly Ser
          20          25          30

Met Met Pro Val Gln Ala Asp Met Pro Arg Pro Thr Gly Leu Ala Ala
          35          40          45

Asp Ile Arg Trp Thr Ala Tyr Gly Val Pro His Ile Arg Ala Lys Asp
          50          55          60

Glu Arg Gly Leu Gly Tyr Gly Ile Gly Tyr Ala Tyr Ala Arg Asp Asn
65          70          75          80

Ala Cys Leu Leu Ala Glu Glu Ile Val Thr Ala Arg Gly Glu Arg Ala
          85          90          95

Arg Tyr Phe Gly Ser Glu Gly Lys Ser Ser Ala Glu Leu Asp Asn Leu
          100          105          110

Pro Ser Asp Ile Phe Tyr Ala Trp Leu Asn Gln Pro Glu Ala Leu Gln
          115          120          125

Ala Phe Trp Gln Ala Gln Thr Pro Ala Val Arg Gln Leu Leu Glu Gly
          130          135          140

Tyr Ala Ala Gly Phe Asn Arg Phe Leu Arg Glu Ala Asp Gly Lys Thr
145          150          155          160

Thr Ser Cys Leu Gly Gln Pro Trp Leu Arg Ala Ile Ala Thr Asp Asp
          165          170          175

Leu Leu Arg Leu Thr Arg Arg Leu Leu Val Glu Gly Gly Val Gly Gln
          180          185          190

Phe Ala Asp Ala Leu Val Ala Ala Ala Pro Pro Gly Ala Glu Lys Val
          195          200          205

Ala Leu Ser Gly Glu Gln Ala Phe Gln Val Ala Glu Gln Arg Arg Gln
          210          215          220

Arg Phe Arg Leu Glu Arg Gly Ser Asn Ala Ile Ala Val Gly Ser Glu
225          230          235          240

```

Arg Ser Ala Asp Gly Lys Gly Met Leu Leu Ala Asn Pro His Phe Pro  
 245 250 255  
 Trp Asn Gly Ala Met Arg Phe Tyr Gln Met His Leu Thr Ile Pro Gly  
 260 265 270  
 Arg Leu Asp Val Met Gly Ala Ser Leu Pro Gly Leu Pro Val Val Asn  
 275 280 285  
 Ile Gly Phe Ser Arg His Leu Ala Trp Thr His Thr Val Asp Thr Ser  
 290 295 300  
 Ser His Phe Thr Leu Tyr Arg Leu Ala Leu Asp Pro Lys Asp Pro Arg  
 305 310 315 320  
 Arg Tyr Leu Val Asp Gly Arg Ser Leu Pro Leu Glu Glu Lys Ser Val  
 325 330 335  
 Ala Ile Glu Val Arg Gly Ala Asp Gly Lys Leu Ser Arg Val Glu His  
 340 345 350  
 Lys Val Tyr Gln Ser Ile Tyr Gly Pro Leu Val Val Trp Pro Gly Lys  
 355 360 365  
 Leu Asp Trp Asn Arg Ser Glu Ala Tyr Ala Leu Arg Asp Ala Asn Leu  
 370 375 380  
 Glu Asn Thr Arg Val Leu Gln Gln Trp Tyr Ser Ile Asn Gln Ala Ser  
 385 390 395 400  
 Asp Val Ala Asp Leu Arg Arg Arg Val Glu Ala Leu Gln Gly Ile Pro  
 405 410 415  
 Trp Val Asn Thr Leu Ala Ala Asp Glu Gln Gly Asn Ala Leu Tyr Met  
 420 425 430  
 Asn Gln Ser Val Val Pro Tyr Leu Lys Pro Glu Leu Ile Pro Ala Cys  
 435 440 445  
 Ala Ile Pro Gln Leu Val Ala Glu Gly Leu Pro Ala Leu Gln Gly Gln  
 450 455 460  
 Asp Ser Arg Cys Ala Trp Ser Arg Asp Pro Ala Ala Ala Gln Ala Gly  
 465 470 475 480  
 Ile Thr Pro Ala Ala Gln Leu Pro Val Leu Leu Arg Arg Asp Phe Val  
 485 490 495  
 Gln Asn Ser Asn Asp Ser Ala Trp Leu Thr Asn Pro Ala Ser Pro Leu  
 500 505 510  
 Gln Gly Phe Ser Pro Leu Val Ser Gln Glu Lys Pro Ile Gly Pro Arg  
 515 520 525  
 Ala Arg Tyr Ala Leu Ser Arg Leu Gln Gly Lys Gln Pro Leu Glu Ala  
 530 535 540

Lys Thr Leu Glu Glu Met Val Thr Ala Asn His Val Phe Ser Ala Asp  
 545 550 555 560  
 Gln Val Leu Pro Asp Leu Leu Arg Leu Cys Arg Asp Asn Gln Gly Glu  
 565 570 575  
 Lys Ser Leu Ala Arg Ala Cys Ala Ala Leu Ala Gln Trp Asp Arg Gly  
 580 585 590  
 Ala Asn Leu Asp Ser Gly Ser Gly Phe Val Tyr Phe Gln Arg Phe Met  
 595 600 605  
 Gln Arg Phe Ala Glu Leu Asp Gly Ala Trp Lys Glu Pro Phe Asp Ala  
 610 615 620  
 Gln Arg Pro Leu Asp Thr Pro Gln Gly Ile Ala Leu Asp Arg Pro Gln  
 625 630 635 640  
 Val Ala Thr Gln Val Arg Gln Ala Leu Ala Asp Ala Ala Ala Glu Val  
 645 650 655  
 Glu Lys Ser Gly Ile Pro Asp Gly Ala Arg Trp Gly Asp Leu Gln Val  
 660 665 670  
 Ser Thr Arg Gly Gln Glu Arg Ile Ala Ile Pro Gly Gly Asp Gly His  
 675 680 685  
 Phe Gly Val Tyr Asn Ala Ile Gln Ser Val Arg Lys Gly Asp His Leu  
 690 695 700  
 Glu Val Val Gly Gly Thr Ser Tyr Ile Gln Leu Val Thr Phe Pro Glu  
 705 710 715 720  
 Glu Gly Pro Lys Ala Arg Gly Leu Leu Ala Phe Ser Gln Ser Ser Asp  
 725 730 735  
 Pro Arg Ser Pro His Tyr Arg Asp Gln Thr Glu Leu Phe Ser Arg Gln  
 740 745 750  
 Gln Trp Gln Thr Leu Pro Phe Ser Asp Arg Gln Ile Asp Ala Asp Pro  
 755 760 765  
 Gln Leu Gln Arg Leu Ser Ile Arg Glu  
 770 775

<210> 6

<211> 33

<212> DNA

<213> Artificial Sequence

<220>

<223> Forward Primer for the QsbA gene

<400> 6

cgtggatcca tgatgcagga ttcgccgctg cgc

33

<210> 7

<211> 32

<212> DNA

<213> Artificial Sequence

<220>

<223> Reverse Primer for the QsbA gene

<400> 7

cgcgattca ccggcagccc tcatgcgaca ac

32

## INTERNATIONAL SEARCH REPORT

International Application No.

PCT/SG 000011

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N9/80 C12N15/57 A01K67/027 A61K38/50

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the International search (name of data base and, where practical, search terms used)

EPO-Internal, BIOSIS, EMBL, WPI Data

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>DATABASE SWISS-PROT 'Online! EBI-SBI; 1 May 2000 (2000-05-01) WHITE ET AL.: "Genome sequence of the radioresistant bacterium Deinococcus radiodurans R1; Science 286:1571-1577 (1999)" retrieved from SWISS-PROT Database accession no. Q9RYQ4 XP002208348 "Aculeacin A acylase from Deinococcus radiodurans R1; 53,316% identity with SEQ ID NO: 2 in 769 aa overlap". abstract</p> <p style="text-align: center;">--- -/-</p>	4-6

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

## \* Special categories of cited documents:

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the International filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the International filing date but later than the priority date claimed

- \*T\* later document published after the International filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- \* & \* document member of the same patent family

Date of the actual completion of the international search

14 August 2002

Date of mailing of the international search report

02/09/2002

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Ury, A



## INTERNATIONAL SEARCH REPORT

 International Application No.  
 PCT/SG 00011

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	DATABASE EMBL 'Online! EBI; 23 November 1999 (1999-11-23) WHITE ET AL.: "Genome sequence of the radioresistant bacterium Deinococcus radiodurans R1; Science 286:1571-1577 (1999)" Database accession no. AE001836 XP002208349 "CDS complement (53991..56348): Encoding Aculeacin A acylase from Deinococcus radiodurans R1: 62,025% identity with SEQ ID NO: 1 in 1817 nt overlap". ---	2
X	WO 01 98214 A (NOVOZYMES BIOTECH INC) 27 December 2001 (2001-12-27) the whole document	2,4-6, 10-20
Y	same citations	2,4-6, 10-20
Y	--- LEADBETTER JARED R ET AL: "Metabolism of acyl-homoserine lactone quorum-sensing signals by Variovorax paradoxus." JOURNAL OF BACTERIOLOGY, vol. 182, no. 24, December 2000 (2000-12), pages 6921-6926, XP002208346 ISSN: 0021-9193 cited in the application abstract; figure 7 ---	2,4-6, 10,20
Y	LEADBETTER JARED R: "Quieting the raucous crowd" NATURE, vol. 411, 14 June 2001 (2001-06-14), pages 748-749, XP002208347 Figure 7 the whole document ---	2,4-6, 10-20
A	--- DONG YI-HU ET AL: "AiiA, an enzyme that inactivates the acylhomoserine lactone quorum-sensing signal and attenuates the virulence of Erwinia carotovora" PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF USA, NATIONAL ACADEMY OF SCIENCE. WASHINGTON, US, vol. 97, no. 7, 28 March 2000 (2000-03-28), pages 3526-3531, XP002166712 ISSN: 0027-8424 cited in the application abstract --- -/--	1-20

## INTERNATIONAL SEARCH REPORT

International Application No  
PCT/SG 00011

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>DONG YI-HU ET AL.: "Quenching quorum-sensing-dependent bacterial infection by an N-acyl homoserine lactonase"</p> <p>NATURE, vol. 411, 14 June 2001 (2001-06-14), pages 813-817, XP001093866 cited in the application abstract</p> <p>-----</p>	1-20

International Publication No  
PCT/SG 00011

Form PCT/ISA/210 (patent family annex) (July 1992)

**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☒ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☒ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**